

TENAZOLE 250 EW

Reg. No.: L8558 Act No. 36 of 1947 • Namibia: N-AR1628 • Botswana: W130822

A systemic fungicide emulsion oil, in water, for the control of the diseases mentioned on the crop listed.

FRAC FUNGICIDE GROUP CODE: 3

ACTIVE INGREDIENT:

Tebuconazole (triazole) 250 g/ℓ

REGISTRATION HOLDER:


Arysta LifeScience South Africa (PTY) LTD
Co. Reg. No.: 2009/019713/07
7 Sunbury Office Park , Off Douglas Saunders Drive
La Lucia, 4019 • Tel: (031) 514-5600

DISTRIBUTOR:

PROTEK, a division of PE-BEE Agri (PTY) LTD
Co. Reg. No.: 2005/036308/07
P.O. Box 72, Heidelberg, 1438
Tel: (011) 812 9800 or 0861 PROTEK (0861 77 68 35)
www.proteksa.co.za

EMERGENCY NO.: 082 446 8946

UN No.: 2757



WARNINGS:

Withholding periods. (Minimum number of days between last application and harvest or grazing and/or feeding).

Barley and Wheat	77 days	Beans & Potatoes:	14 days
Oats	56 days	Tomatoes & Peas:	7 days
Mangoes	Do not apply after 100% petal drop	Groundnut and Bean Hay	42 days

- Harmful when absorbed through the skin or if swallowed.
- May cause serious damage to eyes.
- Toxic to fish.
- Store in a cool place away from food and feedstuffs.
- Keep out of reach of children, uninformed persons, and animals.
- Re-entry interval. Do not enter treated area until spray deposit has dried unless wearing protective clothing.

Although this remedy has been intensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions; quality of dilution water; compatibility with other substances not indicated on the label and the occurrence of resistance of the disease against the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or the occurrence of conditions which could not have been foreseen in the terms of the registration. Consult the supplier in the event of any uncertainty

PRECAUTIONS:

- Do not inhale fumes or spray mist.
- Wear a face shield when handling concentrate
- Wash contaminated clothing daily.
- Wash with soap and water after use.
- Do not eat, drink or smoke whilst mixing, applying or before washing hands and face.
- Avoid drift of spray onto other crops, grazing, rivers, dams and areas not under treatment.
- Clean applicator after use, dispose of wash water, where it will not contaminate crops, grazing, rivers, and dams.
- Rinse empty container three times with a volume of water equal to a minimum of 10% of that of the container. Add the rinsings to the spray tank before destroying the empty container by perforation and flattening.
- Never use empty container for any other purpose.
- Prevent contamination of food, feedstuffs, drinking water, and eating utensils.

RESISTANCE WARNING:

For resistance management **TENAZOLE 250 EW** is a group code 3 fungicide (or a DMI fungicide). Any fungus population may contain individuals naturally resistant to **TENAZOLE 250 EW** and other group code 3 fungicides. The resistant individuals can eventually dominate the fungus population if these fungicides are used repeatedly. These resistant fungi may not be controlled by **TENAZOLE 250 EW** or any other group 3 fungicides. To delay fungicide resistance:

- Avoid exclusive repeated use of fungicides from the same fungicide group code. Alternate or tank mix with products from different fungicide codes.
- For tank mixing or alternation with products in fungicide group code M, refer to applicable product labels.
- Integrate control methods (chemical, cultural and biological) into disease control programs.

For specific information on resistance management contact the registration holder of this product.

DIRECTIONS OF USE: Use only as directed.

COMPATIBILITY:

TENAZOLE 250 EW is compatible with most insecticides and fungicides. The compatibility of **TENAZOLE 250 EW** may be influenced by the formulations of the other products as well as the quality of the spray water. Since the formulations of products may change without the knowledge of Arysta LifeScience and water quality may differ from farm to farm it is advisable to carry out a physical compatibility test before spraying is commenced. Please contact your representative for more details. When tank mixtures are made with grass herbicides to be applied to cereals slight leaf mottling/scorch may occur but new growth will not be affected.

MIXING INSTRUCTIONS:

Fill the spray tank half full with water and pour the required quantity of **TENAZOLE 250 EW** into the water while agitating. Thereafter fill the tank with water. Continue agitation during filling and spraying. Do not allow the **TENAZOLE 250 EW** spray mixture to stand any length of time before using.

GROUND APPLICATION:

TENAZOLE 250 EW can be applied with a conventional high pressure sprayer that is equipped with an effective agitator. It should be equipped with hollow cone nozzles to deliver medium to fine droplets. The sprayer must be correctly calibrated to ensure that the spray mixture is applied uniformly over the target area.

CROP/DISEASE	DOSAGE RATE	REMARKS
BARLEY: Leaf spot (<i>Rhynchosporium secalis</i>)	15 mℓ/10 ℓ water	Apply when first symptoms are noticed or between the seven-leaf and flag leaf stages. A single application should be sufficient for control.
Leaf rust (<i>Puccinia hordei</i>)	25 mℓ/10 ℓ water	Spray as soon as first symptoms appear. Repeat the application after four weeks should the symptoms reappear.
Powdery mildew (<i>Erysiphe graminis</i>)	25 mℓ/10 ℓ water	Spray as soon as first symptoms appear. Repeat the application after four weeks should the symptoms reappear.
Net blotch (<i>Pyrenophora teres</i>) (<i>Helminthosporium</i>)	25 mℓ/10 ℓ water	Apply when first symptoms are noticed especially from the flag leaf stage to ear emergence stage. Under condition of high infection pressure a second application 3-4 weeks should be applied.
Halo spot (<i>Selenophoma donacis</i>)	25 mℓ/10 ℓ water	Net blotch applications will also control Halo Spot.
Eye spot (<i>Pseudocercospora herpotrichoides</i>)	25 mℓ/10 ℓ water	Apply between the seven-leaf and flag leaf stages. A single application should be sufficient for control.

CROP/DISEASE	DOSAGE RATE	REMARKS
WHEAT: Apply TENAZOLE 250 EW at the first signs of infections before the 5% infection level is reached. A second application 21 days after the initial application is recommended for wheat with a high yield potential, particularly where the initial application was made early (first node stage) or when other diseases develop later in the season.		
Eye spot (<i>Pseudocercospora herpotrichoides</i>)	25 mℓ/10 ℓ water	Apply at first to second node stage.
Powdery mildew (<i>Erysiphe graminis</i>)	25 mℓ/10 ℓ water	Spray as soon as the first symptoms are noticed, but not later than the flag leaf stage. Repeat the application should the symptoms reappear.
Speckled leaf blotch (<i>Septoria tritici</i>)	20 mℓ/10 ℓ water	Apply when first symptoms are noticed but not later than the appearance of flag leaf.
Yellow (stripe) rust (<i>Puccinia striiformis</i>)	20 mℓ/10 ℓ water	Apply when the first symptoms are noticed. Repeat the application should the symptoms reappear.
Leaf rust (<i>Puccinia recondita</i>)	25 mℓ/10 ℓ water	Spray as soon as the first symptoms are noticed. Repeat the application should the symptoms reappear.
Glume blotch (<i>Septoria nodorum</i>)	25 mℓ/10 ℓ water	Apply from second node stage but not later than ear emergence.
Stem rust (<i>Puccinia graminis</i>)	30 mℓ/10 ℓ water	Apply at the first signs of stem rust infections, but not later than the flag leaf stage. Repeat the application should symptoms reappear. Effective control of stem rust is dependent on good coverage of the plant and particularly the stem. Ground application is therefore preferable. Aerial application may result in inadequate coverage of stems, especially of densely planted wheat when applied after flag stage. This will result in poor control.
OATS: Crown rust (<i>Puccinia coronata</i>)	15 mℓ/10 ℓ water	Apply as soon as the first sign of infection are noticed.
MANGOES; Powdery mildew (<i>Oidium mangiferae</i>)	20 mℓ/100 ℓ water	Commence application at the first signs of infection, usually when 50 % of the flowers are open. Repeat at intervals of 10 – 14 days up to 100 % petal drop (2 – 3 sprays depending on weather conditions and uniformity of the flowering period). Apply as a cover spray. Ensure thorough coverage of the flower panicles.
POTATOES: Early blight (<i>Alermaria solani</i>)	7.5 mℓ/10 ℓ water	Apply TENAZOLE 250 EW preventatively or apply the first spray at the first signs of the disease. Apply the follow-up sprays at 7 to 10 day intervals. Apply the higher dosage rate (aerial application) after the flowering stage or with warm, moist climatic conditions which favour development of Early blight. If conditions favour the development of Late blight, apply a tank mixture of TEBUZOLE 250 EW with a registered Late blight product. Alternatively TENAZOLE 250 EW sprays can, as necessary, be alternated each 7 to 10 days with a different chemical group fungicide, which is registered for Early and/or Late blight. Do not apply more than 5 TENAZOLE 250 EW sprays per season.
TOMATOES: Early blight (<i>Alermaria solani</i>)	7.5 mℓ/10 ℓ water	Apply 5 – 12 ℓ spray mixture per 100m row length depending on the plant height. Ensure thorough coverage of the plants. Commence application at the first signs of infection. TENAZOLE 250 EW should be applied in a tank mixture with a registered late blight fungicide when conditions are favourable for the development of late blight. TENAZOLE 250 EW should not be applied more than 5 times per season. Apply every 7 – 10 days or alternate every 7– 10 days in a program with a treatment (non-DMI fungicide) registered against early and/or late blight, depending on conditions.
GROUNDNUTS: Leaf spots (<i>Cercospora arachidicola</i> & <i>Cercosporidium personatum</i>), Leaf blotch (<i>Phoma arachidicola</i>) Rust (<i>Puccinia arachidis</i>)	8 mℓ/10 ℓ water 12 mℓ/10 ℓ water	DRY LAND: Apply TENAZOLE 250 EW preventatively or apply the first spray at the first signs of the diseases. Apply 4 to 5 follow-up sprays per season, at 10 to 14 day intervals. Ensure thorough wetting of the foliage. The addition of a registered wetter is recommended. IRRIGATION: Apply TENAZOLE 250 EW preventatively or apply the first spray at the first signs of the diseases. Apply 4 to 5 follow-up sprays per season, at 10 to 14 day intervals. Ensure thorough wetting of the foliage. The addition of a registered wetter is recommended. Ground application: Apply in 500 litres water per hectare.
ONIONS: Purple blotch (<i>Alternaria porri</i>)	15 mℓ/10 ℓ water	DO NOT APPLY TO SPRING ONIONS. Apply as a full cover application at first signs of disease. Repeat every 7 – 14 days as necessary. Do not apply more than 6 times per season. Add a suitable wetter/sticker to avoid excessive run-off.
BEANS: Rust (<i>Uromyces appendiculatus</i>)	10 mℓ/10 ℓ water	Apply as a full cover, preventive treatment or as soon as the first symptoms of the disease is noticed. Repeat every 10-12 days if necessary.
PEAS: Powdery mildew (<i>Erysiphe pisi</i>)	5 mℓ/10 ℓ water	Apply as a full cover application at first signs of disease. Repeat every 10 – 14 days as necessary. Do not apply more than 3 times per season. Add a suitable wetter/sticker to avoid excessive run-off.
FLOWERS AND ORNAMENTALS (ROSES) Powdery mildew (<i>Sphaerotheca pannose</i>) Black spot (<i>Antinonema rosae</i>)	10 mℓ/10 ℓ water	Apply as a full cover, preventive treatment or as soon as the first symptoms of disease is noticed. Repeat every 7 days if necessary.

TENAZOLE 250 EW

Reg. No.: L8558 Wet No. 36 van 1947 • Namibië: N-AR1628 • Botswana: W130822

'n Sistemiese swamdoder emulsie, olie in water, vir die beheer van siektes op gewasse soos genoem.

FRAC SWAMDODER GROEPKODE: 3

AKTIEWE BESTANDEEL:

Tebukonasool (triasool) 250 g/ℓ

VERSPREIDER:


PROTEK, 'n divisie van **PE-BEE Agri (EDMS) BPK**
Mpy. Reg. Nr.: 2005/036308/07
Posbus 72, Heidelberg, 1438
Tel: (011) 812 9800 of 0861 PROTEK (0861 77 68 35)
www.proteksa.co.za

REGISTRASIEHOUER:

Arysta LifeScience South Africa (EDMS) BPK
Mpy. Reg. Nr.: 2009/019713/07
7 Sunbury Office Park
Douglas Saunders-rylaan
La Lucia, 4019 • Tel: (031) 514-5600

NOOD NOMMER: 082 446 8946

UN Nr.: 2757



WAARSKUWINGS:

Onthoud insperiodes. (Minimum aantal dae tussen toediening en oes of beweiding en/of voer).

Gars en koring	77 dae	Boontjies en Aartappels	14 dae
Hawer	56 dae	Tamaties en Ertjies	7 dae
Mango's	Moenie na 100% blomblaarval toedien nie	Grondbone en Boontjehooi	42 dae

- Skadelik indien ingestuk of deur vel geabsorbear word.
- Mag ernstige oogskade veroorsaak.
- Giftig vir visse.
- Stoor in 'n koel plek weg van voedsel en voer.
- Hou buite bereik van kinders, oningeligte persone en diere.
- **HERBETREDINGSINTERVAL.** Moet nie behandelde gebied binnegaan voordat die spuitneerslag droog is nie, tensy beskermende klere gedra word.

Alhoewel hierdie middel omvattend onder 'n groot verskeidenheid toestande getoets is, waarborg die registrasiehouer nie dat dit onder alle toestande doeltreffend sal wees nie aangesien die werking en effek daarvan beïnvloed kan word deur faktore soos abnormale grond-, klimaats- en bergings toestande; kwaliteit van verdunningswater, verenigbaarheid met ander stowwe wat nie op die etiket aangedui is nie en die voorkoms van weerstand van die siekte teen die betrokke middel, sowel as die metode, tyd en akkuraatheid van toediening. Verder aanvaar die registrasiehouer nie verantwoordelikheid vir skade aan gewasse, plantegroei, die omgewing, of vir nadelige effek op mens of dier of vir 'n gebrek aan prestasie van die betrokke middel as gevolg van die versuim van die gebruiker om etiketaanwysings na te kom of as gevolg van die ontstaan van toestande wat nie kragtens die registrasie voorsien kon word nie. Raadpleeg die verskaffer in die geval van enige onsekerheid.

VOORSORGMATREELS:

- Moet nie dampe of spuitnewel inasem nie.
- Dra 'n gesigskerm wanneer konsentraat hanteer word.
- Was besoedelde klere daaglik.
- Was met seep en water na gebruik.
- Moet nie eet, drink of rook tydens vermenging, toediening of voordat hande en gesig gewas is nie.
- Vermoegdrywing van spuitnewel na ander gewasse, weiding, riviere, damme en gebiede wat nie behandel word nie.
- Maak toedieningsapparaat skoon na gebruik en gooi waswater weg waar dit nie gewasse, weiding, riviere en damme sal besoedel nie.
- Spoel lee houder driekeer uit met 'n volume water gelykstaande aan 'n minimum van 10% van die houder. Gooi die speelwater by die inhoud van die spuittenk voordat die houder vernietig word deur plat te slaan en gate in te slaan.
- Moet leë houder nooit vir enige ander doeleindes gebruik nie.
- Voorkom besoedeling van voedsel, voer, drinkwater en eetgerei.

BESTANDEIDSWAARSKUWING:

TENAZOLE 250 EW 'n groepkode 3 swamdoder (of 'n DMI swamdoder). Enige populasie van 'n spesifieke swam mag individue insluit wat 'n natuurlike weerstand teen **TENAZOLE 250 EW** of enige ander groepkode 3 swamdoder het. Indien hierdie swamdoders herhaaldelik aangewend word, kan die weerstandbiedende individue uiteindelik die swampopulasie oorheers. Hierdie weerstandbiedende swamme sal waarskynlik nie deur **TENAZOLE 250 EW** of enige ander groepkode 3 swamdoder beheer word nie. Om weerstand teen swamdoders te vertraag:

- Vermoeg die eksklusiewe herhaaldelike gebruik van swamdoders met dieselfde groepkode. Wissel af met of gebruik tenkmengsels van produkte in verskillende swamdoder groepkodes.
- Raadpleeg toepaslike individuele produketikette wanneer daar afgewissel word, of tenkmengsels gemaak word, met produkte in swamdoder groepkode M.
- Integreer ander beheermaatreels (chemies, verbouing en biologies) in swamdoder programme.

Vir spesifieke inligting oor weerstandsbestuur kontak die registrasiehouer van hierdie produk.

GEBRUIKSAANWYSINGS: Gebruik slegs soos aangedui.

VERENIGBAARHEID:

TENAZOLE 250 EW is mengbaar met meeste insekdoders en swamdoders. Die verenigbaarheid van **TENAZOLE 250 EW** met ander produkte kan beïnvloed word deur die formulering van die betrokke produkte asook die kwaliteit van die verdunningswater. Aangesien die formulering van produkte kan verander sonder die medewete van Arysta LifeScience en die waterkwaliteit van plaas tot plaas kan verskil, moet 'n fisiese verenigbaarheidstoets altyd uitgevoer word voordat bespuiting begin. Kontak u verteenwoordiger vir meer besonderhede. Wanneer tenkmengsels met graskonruiddoders gemaak word om aan graangewasse toe te dien kan ligte spikkels of blaarskroei voorkom. Die nuwe groei sal egter nie nadelig beïnvloed word nie.

MENGINGSTRUKSIES:

Maak die spuittenk halfvol met water en voeg die benodigde hoeveelhede **TENAZOLE 250 EW** by terwyl geroer word. Vul daarna die tenk met water. Dit is belangrik om daarop te let dat die spuitmengsel gedurende vulling en toediening aanhoudend geroer moet word. Moet nie die **TENAZOLE 250 EW** spuitmengsel vir 'n lang tyd laat staan voordat dit gebruik word nie.

GRONDOEDINIENING:

TENAZOLE 250 EW kan met 'n konvensionele hoëvolume spuitapparaat wat toegerus is met 'n doeltreffende roermeganisme, toegedien word. Die spuitstukke moet toegerus wees met holkeëspuitpunte wat medium tot fyn druppels lewer. Om te verseker dat die spuitmengsel eweredig oor die teikengebied versprei word, moet die spuitapparaat korrek gekalibreer wees.

GEWAS / SIEKTE	DOSIS	OPMERKINGS
GARS: Blaarvlek (<i>Rhynchosporium secalis</i>)	15 mL/10 ℓ water	Dien toe wanneer die eerste simptome waargeneem word of tussen die seweblaar- en vlagbaarstadium. 'n Enkele toediening behoort genoegsaam te wees vir beheer.
Blaarroes (<i>Puccinia hordei</i>)	25 mL/10 ℓ water	Dien toe wanneer die eerste simptome waargeneem word. Herhaal toediening na vier weke indien simptome weer voorkom.
Poeieragtige skimmel (<i>Erysiphe graminis</i>)	25 mL/10 ℓ water	Dien toe wanneer die eerste simptome waargeneem word. Herhaal toediening na vier weke indien simptome weer voorkom
Netvlek (<i>Pyrenophora teres</i>) (<i>Helminthosporium</i>)	25 mL/10 ℓ water	Dien toe wanneer die eerste simptome waargeneem word veral vanaf vlagblaarstadium tot aarverskyning. Indien infeksievlekke hoog is, herhaal 3 – 4 weke later.
Ringvlek (<i>Selenophoma donacis</i>)	25 mL/10 ℓ water	Netvlek toedienings sal ook ringvlek beheer.
Oogvlek (<i>Pseudocercospora herpotrichoides</i>)	25 mL/10 ℓ water	Dien toe vanaf die seweblaar- en vlagblaarstadium. 'n Enkele toediening behoort genoegsaam te wees vir beheer.

GEWAS / SIEKTE	DOSIS	OPMERKINGS
KORING: Dien TENAZOLE 250 EW toe by die eerste tekens van besmetting voordat die 5% infeksievlek bereik is. 'n Tweede toediening 21 dae na die eerste toediening word aanbeveel vir koring met 'n hoë opbrengs potensiaal, veral waar die eerste toediening vroeg toegedien is (eerste nodestadium) of waar ander siektes later in die seisoen ontwikkel.		
Oogvlek (<i>Pseudocercospora herpotrichoides</i>)	25 mL/10 ℓ water	Dien toe by die eerste tot tweede nodestadium.
Poeieragtige skimmel (<i>Erysiphe graminis</i>)	25 mL/10 ℓ water	Dien toe wanneer die eerste simptome waargeneem word maar nie later as die vlagblaarstadium nie. Herhaal indien die simptome weer verskyn.
Vaalblaar (<i>Septoria tritici</i>)	20 mL/10 ℓ water	Dien toe wanneer die eerste simptome waargeneem word maar nie later as die vlagblaarstadium nie.
Streeproes (Geelroes) (<i>Puccinia striiformis</i>)	20 mL/10 ℓ water	Dien toe wanneer die eerste simptome waargeneem word maar nie later as die vlagblaarstadium nie.
Blaarroes (<i>Puccinia recondita</i>)	25 mL/10 ℓ water	Dien toe wanneer die eerste simptome waargeneem word. Herhaal indien die simptome weer verskyn.
Bruinaarsiekte (<i>Septoria nodorum</i>)	25 mL/10 ℓ water	Dien toe vanaf die tweede nodestadium maar nie later as aarverskyning nie.
Stamroes (<i>Puccinia graminis</i>)	30 mL/10 ℓ water	Dien toe wanneer die eerste simptome waargeneem word maar nie later as die vlagblaarstadium nie. Herhaal indien die simptome weer verskyn. Effektiewe beheer van stamroes is onderhewig aan goeie bedekking van die plant, veral die stam. Grondtoediening is gevolglik verkieslik. Lugtoediening mag lei tot onvoldoende bedekking van die stamme, veral in dig geplante koring wanneer toediening na die vlagblaarstadium plaasvind. Beheer sal dan swak wees.
HAWER: Kroonroes (<i>Puccinia coronata</i>)	15 mL/10 ℓ water	Dien toe wanneer die eerste simptome waargeneem word.
MANGO'S: Poeieragtige skimmel (<i>Oidium mangiferae</i>)	20 mL/100 ℓ water	Dien toe wanneer die eerste simptome opgemerk word, gewoonlik wanneer 50% van die blomme oop is. Herhaal elke 10 – 14 dae tot en met 100% blomblaarval (2 – 3 bespuitings, afhange van weertoestande en uniformiteit van die blomperiode). Dien toe as 'n dekttoediening en verseker dat die blomtrosse deeglik benat is.
AARTAPPELS: Vroeë roes (<i>Alemaria solani</i>)	7.5 mL/10 ℓ water	Dien TENAZOLE 250 EW voorkomend of dien die eerste bespuiting toe met die eerste tekens van die siekte. Dien die opvolgbespuitings toe met 7 tot 10 dae intervalle. Gebruik die hoër dosis (lugtoediening) na die blomstadium of met warm, klam klimaatstoestande wat gunstig is vir ontwikkeling van vroeë roes. Indien toestande gunstig is vir die ontwikkeling van laatroes moet 'n tenkmengsel van TENAZOLE 250 EW met 'n geregistreerde laatroes produk gemaak word. Alternatiewelik TENAZOLE 250 EW bespuitings kan, indien nodig, afgewissel word elke 7 tot 10 dae met 'n ander chemiese groep swamdoder, wat geregistreer is vir vroeë en/of laatroes. Moenie meer as 5 TENAZOLE 250 EW bespuitings per seisoen doen nie.
TAMATIES: Vroeë roes (<i>Alemaria solani</i>)	7.5 mL/10 ℓ water	Dien 5 – 12 ℓ spuitmengsel per 100 m ² ry lengte aftangend van die plant hoogte. Verseker deeglike bedekking van die plante. Begin toediening sodra die eerste tekens van infeksie waargeneem word. TENAZOLE 250 EW moet toegedien word in 'n tenkmengsel met 'n geregistreerde laatroes swamdoder wanneer toestande gunstig is vir die ontwikkeling van laatroes. TENAZOLE 250 EW moet nie meer as 5 keer per seisoen toegedien word nie. Dien elke 7 – 10 dae of alternatiewe elke 7 – 10 dae in 'n program met 'n behandeling (nie-DMI swamdoders) wat teen die vroeë en/of laatroes, afhange van omstandighede.
GRONDBONE: Blaarvleksiectes (<i>Cercospora arachidicola</i> & <i>Cercosporidium personatum</i>), Spatselvlek (<i>Phoma arachidicola</i>) Bruinroes (<i>Puccinia arachidis</i>)	8 mL/10 ℓ water 12 mL/10 ℓ water	DROË LAND: Dien TENAZOLE 250 EW voorkomend toe of die eerste bespuiting met die eerste tekens van die siektes. Dien 4 tot 5 opvolgbespuitings per seisoen, op 10 tot 14 dae intervalle. Verseker deeglike bedekking van die blare. Dien byvoeging van 'n geregistreerde benattingsmiddel word aanbeveel. BESPROEÏING: Dien TENAZOLE 250 EW voorkomend toe op die eerste bespuiting met die eerste tekens van die siektes. Dien 4 tot 5 opvolgbespuitings per seisoen, op 10 tot 14 dae intervalle. Verseker deeglike bedekking van die blare. Dien byvoeging van 'n geregistreerde benattingsmiddel word aanbeveel. Grondtoediening: Dien toe in 500 liter water per hektaar.
UIE: Alternaria-vlek (<i>Alternaria porri</i>)	15 mL/10 ℓ water	MOENIE OP SPRIETUIE TOEDIEN NIE. Dien as 'n voldekbepuiting toe met die eerste tekens van die siekte. Herhaal elke 7 – 14 dae as dit nodig is. Moet nie meer as 6 keer per seisoen toedien nie. Voeg 'n geskikte benatter/kleefmiddel by om oormatige afloop te verhoed.
BONE: Bruinroes (<i>Uromyces appendiculatus</i>)	10 mL/10 ℓ water	Dien as 'n voldek, voorkomende behandeling toe of sodra die eerste simptome van die siekte opgemerk word. Herhaal elke 10 – 12 dae indien nodig.
ERTJIES: Poeieragtige skimmel (<i>Erysiphe pisi</i>)	5 mL/10 ℓ water	Dien as 'n voldekbepuiting toe met die eerste tekens van die siekte. Herhaal elke 10 – 14 dae indien nodig. Moet nie meer as 3 keer per seisoen toedien nie. Voeg 'n geskikte benatter/kleefmiddel by om oormatige afloop te verhoed.
BLOMME EN SIERPLANTE (ROSE) Poeieragtige skimmel (<i>Sphaerotheca pannose</i>) Swart vlek (<i>Antinonema rosae</i>)	10 mL/10 ℓ water	Dien as 'n voldek, voorkomende behandeling toe of sodra die eerste simptome van die siekte opgemerk word. herhaal elke 7 dae indien nodig.