

**SCIENTIFIC™
SUPA-KILL**

Supa-Kill

Block Bait

Reg. No. L10422 Act No. 36 of 1947

**READ THE LABEL CAREFULLY AND FOLLOW ALL INSTRUCTIONS
KEEP OUT OF REACH OF CHILDREN AND ANIMALS**

GROUP		RODENTICIDE
--------------	--	--------------------

A HIGHLY ACTIVE ANTICOAGULANT BAIT FOR THE CONTROL OF NORWAY RAT, ROOF RAT, AND HOUSE MICE.

Active Ingredient

Bromadiolone 0.05g/kg



WARNING



Hazard statements:

May be harmful if swallowed.
May be harmful in contact with skin.

Precautionary statements:

If medical advice is needed, have product container or label at hand.
Wear protective gloves.

If medical advice is needed, have product container or label at hand.

NET VOLUME

REGISTERED AND MANUFACTURED BY

Scientific Supa-Kill C.C. CK. No. 1991/027877/23

PO Box 8020, Edleen, 1625

Tel. 011 394 3941/2 Office hours

www.supa-kill.com

EMERGENCY NUMBER

Griffin poison centre: 082 446 8946

Tygerberg poison centre: 086 155 5777

Batch No.:

Date of Manufacture:

UN No.: 3027

Risk and Safety Information

To avoid risk to human health and the environment, comply with the instructions for use.

Hazard statement

- H303: May be harmful if swallowed.
- H313: May be harmful in contact with skin.
- H332: May be harmful if swallowed.

Disclaimer

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because action and effect thereof may be affected by factors such as abnormal climatic and storage conditions; compatibility with other substances not indicated on the label and the occurrence of resistance of the pest against the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

Precautionary statement

- P101: If medical advice is needed, have product container or label at hand.
- P102: Keep out of reach of children.
- P103: Read carefully and follow instructions.

Prevention statement

- P270: Do not eat, drink, or smoke when using this product.
- P280: Wear protective gloves.

WARNINGS

- HANDLE WITH EXTREME CARE.
- Prevent access to bait by children and animals.
- Store away from food and feed.
- Poisonous if swallowed and on prolonged contact with skin.
- Toxic to fish and wildlife.
- In case of poisoning, call a doctor or a veterinarian and make this label available to him.

PRECAUTIONS

- Avoid contamination of feed, food, eating utensils and water.
- Set bait stations where they will be inaccessible to children, pets, livestock, and other non-target animals.
- Wash hands and exposed skin after handling the poison.
- Do not eat, drink or smoke while handling the poison.
- Destroy empty container and do not use for any other purpose.
- Bury dead rodents and any uneaten bait at the end of treatment.

SYMPTOMS OF HUMAN POISONING

- Bleeding from the gums and nose, bleeding under the skin, pallor, haemorrhage into and around the elbow and knee joints. Blood in the urine and faeces.

FIRST AID TREATMENT

- If swallowed, call the Griffon Poison Information Centre (082 446 8946) for treatment advice. Only induce vomiting if advised to do so or if it is less than three hours since the bait was eaten. Take the person to a hospital or a doctor IMMEDIATELY and show the product label. In the case of animal poisonings consult a veterinarian.

NOTE TO PHYSICIAN OR VETERINARIAN

- Bromadiolone is a delayed-action anticoagulant. Perform stomach wash if a large amount of bait was ingested and less than 4 hours have elapsed. Determine prothrombin time 12 to 18 hours after ingestion and if elevated, give Vitamin K1 intravenously in divided doses. In severe cases transfusions of fresh blood may be necessary. CONTACT A POISON INFORMATION CENTRE.

Relevant Compounds

Compound	Weight %
Bromadiolone	0.005
Bitrex	0.001

DIRECTIONS FOR USE

USE ONLY AS DIRECTED

For use in the home, on the farm and industrial premises. Supa-Kill Rat and Mouse poison can kill rodents after a single feed. Death will take place within 4 to 12 days. EFFECTIVE AGAINST WARFARIN RESISTANT RATS AND MICE. ALWAYS PLACE SUPA-KILL RAT & MOUSE POISON IN A COVERED BAIT STATION TO PREVENT ACCESS TO BAIT BY CHILDREN AND ANIMALS. Bait points should be inspected and replenished every 7 days until feeding ceases.

NORWAY AND ROOF RATS

Place 3-4 Supa-Kill Blocks in Supa-Kill bait stations, along walls and active rat runs and in sheltered areas where rats are known to live or feed.

MICE

Place 1-2 Supa-Kill Block's in bait stations every 2 to 4 meters along walls or runways, in nests and in sheltered areas where mice are known to live or feed. The single feeding action of Supa-Kill Rat and Mouse poison is particularly effective against mice with their small appetites and sporadic feeding behaviour.

EMERGENCY NUMBERS

Griffin poison centre: 028 446 8946

Tygerberg poison centre: 086 155 5777